

## Addition of *n*-Butyllithium to an Aldimine: Role of Chelation, Aggregation, and Cooperative Solvation

Bo Qu and David B. Collum\*


Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853-1301

Received March 29, 2005; E-mail: dbc6@cornell.edu

*N,N,N',N'*-Tetramethylethylenediamine (TMEDA) has played a central role in organolithium chemistry.<sup>1</sup> Although its rise to prominence occurred in the context of anionic polymerizations,<sup>2</sup> synthetic organic chemists were quick to exploit TMEDA to accelerate and control a wide range of reactions.<sup>3</sup> TMEDA has also profoundly shaped conventional wisdom about how solvation influences aggregation and reactivity.


Approximately a decade ago we published a polemical review suggesting that TMEDA may not be a universally strong ligand for lithium and that much of the folklore surrounding TMEDA may be ill founded.<sup>4</sup> Among a large number of assertions, we suggested that both competitive and cooperative solvation in TMEDA/THF mixtures might be important, but the data were too meager for definitive conclusions. In passing, we implied that TMEDA/Et<sub>2</sub>O mixtures were much less likely to present such complications.

We describe herein the investigations of 1,2-additions of *n*-BuLi to functionalized imines shown in eq 1.<sup>5–7</sup> The putative advantages offered by TMEDA are challenged at the outset, given that TMEDA attenuates the rates and stereoselectivities. Rate studies reveal a complex mechanistic scenario in which *four* independent mechanisms are detected. Cooperative solvation by Et<sub>2</sub>O and TMEDA is prominent in both monomer- and dimer-based pathways.




*n*-BuLi in TMEDA/toluene is exclusively [(*n*-BuLi)<sub>2</sub>(η<sup>2</sup>-TMEDA)<sub>2</sub>] (**4**) at all accessible *n*-BuLi and TMEDA concentrations.<sup>8,9</sup> The rates of the 1,2-addition of *n*-BuLi<sup>10</sup> to imine **1**<sup>5</sup> (eq 1) were investigated by monitoring the loss of **1** (exclusively uncomplexed; 1667 cm<sup>-1</sup>) using in situ IR spectroscopy<sup>11</sup> as described for analogous *N*-isopropylaldimines.<sup>8b</sup> Approximately 1000-fold higher rates for **1** when compared to the isostructural *N*-isopropyl analogues in TMEDA/toluene (as well as in all TMEDA/Et<sub>2</sub>O combinations described below) attest to the importance of chelation.

A plot of *k*<sub>obsd</sub> versus TMEDA concentration<sup>12</sup> displays an inverse-first-order dependence with a nonzero asymptotic limit (*k*<sub>obsd</sub> = *k*'[TMEDA]<sup>-1.0±0.1</sup> + *k*''[TMEDA]<sup>0</sup>) characteristic of parallel dissociative and nondissociative pathways (Figure 1 inset).<sup>13</sup> Plots of *k*<sub>obsd</sub> versus *n*-BuLi concentration<sup>12</sup> at low and high TMEDA concentrations reveal first-order and half-order dependencies, respectively. Thus, the 1,2-addition is dominated by a dimer-based pathway—[(*n*-BuLi)<sub>2</sub>(TMEDA)(**1**)]<sup>‡</sup>—at low TMEDA concentration and the more sluggish monomer-based pathway—[(*n*-BuLi)-




**Figure 1.** Plot of *k*<sub>obsd</sub> vs TMEDA concentration for the 1,2-addition of *n*-BuLi (0.10 M) to imine **1** (0.007 M) in toluene cosolvent at  $-78\text{ }^{\circ}\text{C}$  with variable Et<sub>2</sub>O concentrations (▽, 6.0 M Et<sub>2</sub>O; ▼, 4.0 M Et<sub>2</sub>O; ○, 2.0 M Et<sub>2</sub>O; ●, 0.0 M Et<sub>2</sub>O). The curves depict unweighted least-squares fits to *k*<sub>obsd</sub> = *a*[TMEDA]<sup>*n*</sup> + *b*. The inset shows the dependence of *k*<sub>obsd</sub> on TMEDA concentration (no Et<sub>2</sub>O) at  $-55\text{ }^{\circ}\text{C}$ .

(TMEDA)(**1**)<sup>‡</sup>—at high TMEDA concentration. We offer transition structures **5** and **6** as reasonable depictions, although we could depict **5** as an open dimer instead (vide infra).<sup>16,17</sup> The five-coordinate lithium of **6** may seem strange at first glance, but evidence of high-coordinate lithium has been accruing.<sup>14,15</sup>




Analogous rate studies in TMEDA/Et<sub>2</sub>O/toluene mixtures afford surprising results. Plots of *k*<sub>obsd</sub> versus TMEDA concentration at several Et<sub>2</sub>O concentrations<sup>18</sup> display inverse-first- and zeroth-order dependencies (Figure 1), suggesting that dimer- and monomer-based pathways are still operative. Indeed, approximate first-order and half-order dependencies on the *n*-BuLi concentration are measured at low and high TMEDA concentrations, respectively. Nonetheless, the rates are markedly sensitive to the Et<sub>2</sub>O concentration, as evidenced in Figure 1. Reprocessing the data in Figure 1 as plots of *k*<sub>obsd</sub> versus Et<sub>2</sub>O concentration (Figure 2) shows that the addition is first-order in Et<sub>2</sub>O concentration at both low and high TMEDA concentrations. Thus, the rate data implicate the dominance by [(*n*-



**Figure 2.** Plot of  $k_{\text{obsd}}$  vs  $\text{Et}_2\text{O}$  concentration for the 1,2-addition of  $n\text{-BuLi}$  (0.10 M) to imine **1** (0.007 M) in toluene cosolvent at  $-78^\circ\text{C}$  with TMEDA (●, 0.10 M TMEDA; ○, 0.20 M TMEDA; ▼, 0.40 M TMEDA; ▽, 0.65 M TMEDA; ■, 0.90 M TMEDA). The curves depict unweighted least-squares fits to  $k_{\text{obsd}} = a[\text{Et}_2\text{O}]^n + b$ .

$\text{BuLi}_2(\text{TMEDA})(\text{Et}_2\text{O})(\mathbf{1})^\ddagger$  at low TMEDA concentration and  $[(n\text{-BuLi})(\text{TMEDA})(\text{Et}_2\text{O})(\mathbf{1})]^\ddagger$  at high TMEDA concentration. Given the strong evidence of chelation by the substrate and the stoichiometries defined by the rate law, we propose transition structures **7** and **8**.



Although there are plausible isomers of dimer-based transition structure **7**, the open dimer motif of **7** is consistent with crystallographic, spectroscopic, computational, and kinetic data,<sup>16,17</sup> and the four-coordinate lithiums seem reasonable. By contrast, the congestion accompanying the high-coordinate lithium in **8** is likely to generate some consternation. It is possible that TMEDA ligands are not chelated. Nonetheless, we are not acutely troubled by six-coordinate (octahedral) lithium.<sup>15</sup>

The monomer- and dimer-based 1,2-additions to **1** contrast with analogous additions to simple imines in which exclusively monomer-based pathways are involved.<sup>8b</sup> The most surprising conclusion, however, is that  $\text{Et}_2\text{O}$  and TMEDA function *cooperatively* in the rate-limiting transition structures.<sup>19</sup> Is cooperative solvation common when mixtures of TMEDA and ethereal solvents are used? Our hunch is yes: Preliminary results on the addition of  $n\text{-BuLi}$ /TMEDA to simple  $N$ -isopropylimines show a first-order dependence on  $\text{Et}_2\text{O}$  concentration. The stereoselectivities (**2:3**) affiliated with the four competing pathways are crudely estimated as follows: **5**, 4:1; **6**,

15:1; **7**, 200:1; **8**, 30:1. Clearly,  $\text{Et}_2\text{O}$  markedly enhances the rates and stereoselectivities of both the dimer- and the monomer-based 1,2-additions. One should remain cognizant of this complexity when optimizing and rationalizing solvent-dependent selectivities.

**Acknowledgment.** We thank the National Science Foundation for direct support of this work as well as DuPont Pharmaceuticals, Merck Research Laboratories, Pfizer, Aventis, R. W. Johnson, Boehringer-Ingelheim, and Schering-Plough for indirect support.

**Supporting Information Available:** NMR spectra and rate data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) *Polyamine-Chelated Alkali Metal Compounds*; Langer, A. W., Jr., Ed.; American Chemical Society: Washington, DC, 1974.
- (2) *Anionic Polymerization: Principles and Practical Applications*; Hsieh, H. L.; Quirk, R. P., Eds.; Marcel Dekker: New York, 1996.
- (3) Snieckus, V. *Chem. Rev.* **1990**, *90*, 879. Langer, P.; Freiberg, W. *Chem. Rev.* **2004**, *104*, 4125. Mangelinckx, S.; Giubellina, N.; De Kimpe, N. *Chem. Rev.* **2004**, *104*, 2353.
- (4) Collum, D. B. *Acc. Chem. Res.* **1992**, *25*, 448.
- (5) Betz, J.; Heuschmann, M. *Tetrahedron Lett.* **1995**, *36*, 4043.
- (6) Recent reviews: Denmark, S. E.; Nicaise, O. J.-C. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, Y., Eds.; Springer-Verlag: Heidelberg, 1999; Chapter 26.2. Kobayashi, S.; Ishitani, H. *Chem. Rev.* **1999**, *99*, 1069. Enders, D.; Reinhold, U. *Tetrahedron: Asymmetry* **1997**, *8*, 1895. Bloch, R. *Chem. Rev.* **1998**, *98*, 1407. Iguchi, M.; Yamada, K.-i.; Tomioka, K. In *Topics In Organometallic Chemistry*; Hodgson, D. M., Ed.; Springer: Heidelberg, 1998; pp 37–59.
- (7) The values of  $k_{\text{rel}}$  and **2:3** ratios in eq 1 were measured under pseudo-first-order conditions described in the text at (1) 0.1 M  $n\text{-BuLi}$  in 0.1 M TMEDA/toluene, (2) 0.1 M TMEDA/6.0 M  $\text{Et}_2\text{O}$ /toluene, and (3) neat  $\text{Et}_2\text{O}$ .
- (8) (a) Hoffmann, D.; Collum, D. B. *J. Am. Chem. Soc.* **1998**, *120*, 5810. (b) Rutherford, J. L.; Hoffmann, D.; Collum, D. B. *J. Am. Chem. Soc.* **2002**, *124*, 264.
- (9) For a bibliography of structural studies of  $n\text{-BuLi}$ /TMEDA, see ref 8a.
- (10)  $n\text{-BuLi}$  was recrystallized from hexane: Kottke, T.; Stalke, D. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 580. Also, see ref 8a.
- (11) Rein, A. J.; Donahue, S. M.; Pavlosky, M. A. *Curr. Opin. Drug Discov. Dev.* **2000**, *3*, 734.
- (12)  $n\text{-BuLi}$  concentration refers to the total concentration (normality). TMEDA concentration refers to the concentrations of the free (uncomplexed) ligand.
- (13) Espenson, J. H. In *Chemical Kinetics and Reaction Mechanisms*; McGraw-Hill: New York, 1995; pp 58–62. For several closely related examples, see: Bernstein, M. P.; Collum, D. B. *J. Am. Chem. Soc.* **1993**, *115*, 789. Ramirez, A.; Candler, J.; Bashore, C. G.; Wirtz, M. C.; Coe, J. W.; Collum, D. B. *J. Am. Chem. Soc.* **2004**, *126*, 14700.
- (14) For leading references to high-coordinate lithium, see: Zhao, P.; Condo, A.; Keresztes, I.; Collum, D. B. *J. Am. Chem. Soc.* **2004**, *126*, 3113.
- (15) For example,  $^+\text{Li}(\text{DME})_3$  is a commonly observed octahedral coordination sphere: Bock, H.; Näther, C.; Havlas, Z.; John, A.; Arad, C. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 875.
- (16) For a structurally related triple ion bearing a TMEDA-chelated internal lithium, see: Bildmann, U. J.; Muller, G. *Organometallics* **2001**, *20*, 1689.
- (17) Open dimer-based mechanisms for 1,2-additions of alkyllithiums have been investigated computationally: Kaufmann, E.; Schleyer, P. v. R.; Houk, K. N.; Wu, Y.-D. *J. Am. Chem. Soc.* **1985**, *107*, 5560. Nakamura, E.; Nakamura, M.; Koga, N.; Morokuma, K. *J. Am. Chem. Soc.* **1993**, *115*, 11016. For additional leading references to spectroscopic, crystallographic, and kinetic evidence of open dimers, see: Zhao, P.; Collum, D. B. *J. Am. Chem. Soc.* **2003**, *125*, 14411.
- (18) NMR spectroscopic studies show that the previously characterized  $[(n\text{-BuLi})_2(\eta^2\text{-TMEDA})_2]$  persists over a wide range of TMEDA– $\text{Et}_2\text{O}$  concentrations to the exclusion of TMEDA– $\text{Et}_2\text{O}$  mixed solvated dimers (Supporting Information). For related studies of phenyllithium in TMEDA/etheral solvent mixtures, see: Reich, H. J.; Green, D. P.; Medina, M. A.; Goldenberg, W. S.; Gudmundsson, B. O.; Dykstra, R. R.; Phillips, N. H. *J. Am. Chem. Soc.* **1998**, *120*, 7201.
- (19) For a crystallographically characterized dimer of  $n\text{-BuLi}$  solvated by both sparteine and  $\text{Et}_2\text{O}$ , see: Strohmenn, C.; Strohmenn, K.; Schildbach, D. *J. Am. Chem. Soc.* **2003**, *125*, 13672.

JA0519987